Brain Research, 441 (1988) 408–410
Elsevier

Author Index

Abercrombie, E.D. and Jacobs, B.L., Systemic naloxone administration potentiates locus coeruleus noradrenergic neuronal activity under stressful but not non-stressful conditions, 362
Agid, Y., see Hirsch, E., 139
Akema, T., Hashimoto, R. and Kimura, F., Preoptic injection of VIP, but not of secretin or PHI, inhibits LH and stimulates prolactin secretion in the ovariectomized rat, 367
Alvarez, J., see Inestrosa, N.C., 331
Amri, M. and Car, A., Projections from the medullary swallowing center to the hypoglossal motor nucleus: a neuroanatomical and electrophysiological study in sheep, 119
Araki, M., McGeer, P.L. and Kimura, H., The efferent projections of the rat lateral habenular nucleus revealed by the PHA-L anterograde tracing method, 319
Audet, M.A., see Doucet, G., 233
Austin, M.C. and Kalivas, P.W., The effect of cholinergic stimulation in the nucleus accumbens on locomotor behavior, 209

Bak, T.H., see Carr, K.D., 173
Barman, S.M. and Gebber, G.L., The axons of raphespinal sympathoinhibitory neurons branch in the cervical spinal cord, 371
Benedetti, M., see Frank, C., 393
Berger, B., Doucet, G. and Descarries, L., Density of the dopamine innervation in rat cerebral cortex after neonatal 6-hydroxydopamine or adult stage DSP-4 noradrenaline denervations: a quantitative radioautographic study, 260
Berger, B., see Doucet, G., 233
Black, I.B., see La Gamma, E.F., 292
Brown, M.R., Hauger, R. and Fisher, L.A., Autonomic and cardiovascular effects of corticotropin-releasing factor in the spontaneously hypertensive rat, 33
Buéno, L. and Gué, M., Evidence for the involvement of corticotropin-releasing factor in the gastrointestinal disturbances induced by acoustic and cold stress in mice, 1

Cabib, S., Kempf, E., Schleef, C., Oliverio, A. and Puglisi-Allegra, S., Effects of immobilization stress on dopamine and its metabolites in different brain areas of the mouse: role of genotype and stress duration, 153
Campbell, J.N., see Meyer, R.A., 81
Campbell, J.N., see Meyer, R.A., 168
Car, A., see Amri, M., 119
Carlsen, J. and Heimer, L., The basolateral amygdaloid complex as a cortical-like structure, 377
Carlton, S.M., see Harmann, P.A., 87
Carr, K.D. and Bak, T.H., Medial thalamic injection of opioid agonists: μ-agonist increases while κ-agonist decreases stimulus thresholds for pain and reward, 173
Chan, J.Y.H., see White, S.R., 195
Costa, M.E., see Ojeda, S.R., 339
Cotter, J.R., see Laemle, L.K., 398
Coyer, P.E., Simeone, F.A. and Michele, J.J., Extended latency of the cortical component of the somatosensory-evoked potential accompanying moderate increases in cerebral blood flow during systemic hypoxia in cats, 145

Dardenne, M., see Hirsch, E., 139
Decker, J., see Jenny, A, 352
Descarries, L., see Berger, B., 260
Descarries, L., see Doucet, G., 233
Diadori, P., see Dykes, R.W., 48
Dolly, J.O., see Pelchen-Matthews, A., 127
Doucet, G., Descarries, L., Audet, M.A., Garcia, S. and Berger, B., Radioautographic method for quantifying monoamine innervations in the rat brain. Application to the cerebral cortex, 233
Doucet, G., see Berger, B., 260
Dudley, C.A. and Moss, R.L., Facilitation of lordosis in female rats by CNS-site specific infusions of an LH-RH fragment, Ac-LH-RH-(5–10), 161
Dunkley, P.R., Heath, J.W., Harrison, S.M., Jarvie, P.E., Glenfield, P.J. and Rostas, J.A.P., A rapid Percoll gradient procedure for isolation of synaptosomes directly from an S_1 fraction: homogeneity and morphology of subcellular fractions, 59
Dunkley, P.R., see Harrison, S.M., 72
Dutar, P., see Dykes, R.W., 48
Dykes, R.W., Lamour, Y., Diadori, P., Landry, P. and Dutar, P., Somatosensory cortical neurons with an identifiable electrophysiological signature, 48

Ernsberger, P., Meeley, M.P. and Reis, D.J., An endogenous substance with clonidine-like properties: selective binding to imidazole sites in the ventrolateral medulla, 309

Fisher, L.A., see Brown, M.R., 33
Frank, C., Sagratella, S., Benedetti, M. and Scotti de Carolis, A., Comparative influence of calcium blocker and purinergic drugs on epileptiform bursting in rat hippocampal slices, 393
Freeman, J.A., see Langdon, R.B., 299
Fujimoto, K., see Sakata, T., 403
Fukagawa, K., see Sakata, T., 403
Fukuda, T. and Jacobowitz, D.M., Purification and immunocytochemical detection of a protein that reveals layer V pyramidal cells in the rat cortex, 185

Garcia, S., see Doucet, G., 233
Gebber, G.L., see Barman, S.M., 371
Gerber, G., see Ryu, P.D., 357
Glenfield, P.J., see Dunkley, P.R., 59
Gramlich, C.A., see Stripling, J.S., 281
Greenough, W.T., see Sirevaag, A.M., 386
Gué, M., see Buéno, L., 1
Guilbert, M., see Guillemot, J.-P., 221
Guillemot, J.-P., Lepore, F., Prevost, L., Richer, L. and Guilbert, M., Somatosensory receptive fields of fibres in the rostral corpus callosum of the cat, 221

Hanesworth, J.M., see Wright, J.W., 215
Harding, J.W., see Wright, J.W., 215

Harmann, P.A., Carlton, S.M. and Willis, W.D., Collaterals of spinothalamic tract cells to the periaqueductal gray: a fluorescent double-labeling study in the rat, 87
Harrison, S.M., Jarvie, P.E. and Dunkley, P.R., A rapid Percoll gradient procedure for isolation of synaptosomes directly from an S_1 fraction: viability of subcellular fractions, 72
Harrison, S.M., see Dunkley, P.R., 59
Hashimoto, R., see Akema, T., 367
Hauger, R., see Brown, M.R., 33
Heath, J.W., see Dunkley, P.R., 59
Heimer, L., see Carlsen, J., 377
Hirsch, E., Ruberg, M., Portier, M.-M., Dardenne, M. and Agid, Y., Characterization of two antigens in parkinsonian Lewy bodies, 139
Hong, J.-S., see Lee, P.H.K., 381
Hutchison, J.B., see Schumacher, M., 98
Hutchison, R.E., see Schumacher, M., 98

Inestrosa, N.C. and Alvarez, J., Axons grow in the aging rat but fast transport and acetylcholinesterase content remain unchanged, 331

Jacobowitz, D.M., see Fukuda, T., 185
Jacobs, B.L., see Abercrombie, E.D., 362
Jarvie, P.E., see Dunkley, P.R., 59
Jarvie, P.E., see Harrison, S.M., 72
Jenny, A., Smith, J. and Decker, J., Motor organization of the spinal accessory nerve in the monkey, 352

Kalivas, P.W., see Austin, M.C., 209
Katz, K.H., see Ojeda, S.R., 339
Kempf, E., see Cabib, S., 153
Kimura, F., see Akema, T., 367
Kimura, H., see Araki, M., 319

Laemle, L.K. and Cotter, J.R., VIP-like immunoreactivity in the anterior hypothalamic area of a hibernating bat varies with physiological state, 398
La Gamma, E.F., White, J.D., McKelvy, J.F. and Black, I.B., Second messenger mechanisms governing opiate peptide transmitter regulation in the rat adrenal medulla, 292
Lamour, Y., see Dykes, R.W., 48
Landry, P., see Dykes, R.W., 48
Langdon, R.B., Manis, P.B. and Freeman, J.A., Goldfish retinotectal transmission in vitro: component current sink–source pairs isolated by varying calcium and magnesium levels, 299
Larson, J. and Lynch, G., Role of *N*-methyl-D-aspartate receptors in the induction of synaptic potentiation by burst stimulation patterned after the hippocampal θ-rhythm, 111
Lee, P.H.K., Obie, J. and Hong, J.-S., Intrahippocampal injections of a specific μ-receptor ligand PL017 produce generalized convulsions in rats, 381
Lepore, F., see Guillemot, J.-P., 221
Lynch, G., see Larson, J., 111

Maderspach, K. and Solomonia, R., Glial and neuronal opioid receptors: apparent positive cooperativity observed in intact cultured cells, 41
Manis, P.B., see Langdon, R.B., 299
McGeer, P.L., see Araki, M., 319
McKelvy, J.F., see La Gamma, E.F., 292
Meeley, M.P., see Ernsberger, P., 309
Meshul, C.K. and Seil, F.J., Transplanted astrocytes reduce synaptic density in the neuropil of cerebellar cultures, 23
Meyer, R.A. and Campbell, J.N., A novel electrophysiological technique for locating cutaneous nociceptive and chemospecific receptors, 81
Meyer, R.A., Campbell, J.N. and Raja, S.N., Antidromic nerve stimulation in monkey does not sensitize unmyelinated nociceptors to heat, 168
Michele, J.J., see Coyer, P.E., 145
Moss, R.L., see Dudley, C.A., 161
Murase, K., see Ryu, P.D., 357

Nakazawa, S., see Takahashi, H., 202

Obie, J., see Lee, P.H.K., 381
Ojeda, S.R., Urbanski, H.F., Katz, K.H. and Costa, M.E., Prostaglandin E_2 releases luteinizing hormone-releasing hormone from the female juvenile hypothalamus through a Ca^{2+}-dependent, calmodulin-independent mechanism, 339
Oliverio, A., see Cabib, S., 153
Ookuma, K., see Sakata, T., 403
Ozaki, H.S. and Shimada, M., The fibers which course within the Probst's longitudinal bundle seen in the brain of a congenitally acallosal mouse: a study with the horseradish peroxidase technique, 5

Patneau, D.K., see Stripling, J.S., 281
Pelchen-Matthews, A. and Dolly, J.O., Distribution of acceptors for β-bungarotoxin in the central nervous system of the rat, 127
Penner, J.D., see White, S.R., 195
Portier, M.-M., see Hirsch, E., 139
Prevost, L., see Guillemot, J.-P., 221
Puglisi-Allegra, S., see Cabib, S., 153

Quirk, W.S., see Wright, J.W., 215

Raja, S.N., see Meyer, R.A., 168
Randic, M., see Ryu, P.D., 357
Reis, D.J., see Ernsberger, P., 309
Richer, L., see Guillemot, J.-P., 221
Rostas, J.A.P., see Dunkley, P.R., 59
Ruberg, M., see Hirsch, E., 139
Ryu, P.D., Gerber, G., Murase, K. and Randic, M., Actions of calcitonin gene-related peptide on rat spinal dorsal horn neurons, 357

Saederup, E., see Squires, R.F., 15
Sagratella, S., see Frank, C., 393
Sakata, T., Ookuma, K., Fukagawa, K., Fujimoto, K., Yoshimatsu, H., Shiraishi, T. and Wada, H., Blockade of the histamine H_1-receptor in the rat ventromedial hypothalamus and feeding elicitation, 403
Schleef, C., see Cabib, S., 153
Schumacher, M., Hutchison, R.E. and Hutchison, J.B., Ontogeny of testosterone-inducible brain aromatase activity, 98
Scotti de Carolis, A., see Frank, C., 393
Seil, F.J., see Meshul, C.K., 23
Shimada, M., see Ozaki, H.S., 5
Shimura, T., see Takahashi, H., 202
Shiraishi, T., see Sakata, T., 403
Simeone, F.A., see Coyer, P.E., 145
Sirevaag, A.M. and Greenough, W.T., A multivariate statistical summary of synaptic plasticity measures in rats exposed to complex, social and individual environments, 386

Smith, J., see Jenny, A., 352
Solomonia, R., see Maderspach, K., 41
Speth, R.C., see White, S.R., 195
Squires, R.F. and Saederup, E., Antidepressants and metabolites that block $GABA_A$ receptors coupled to ^{35}S-*t*-butylbicyclophosphorothionate binding sites in rat brain, 15
Stripling, J.S., Patneau, D.K. and Gramlich, C.A., Selective long-term potentiation in the pyriform cortex, 281

Takahashi, H., Nakazawa, S., Yoshino, Y. and Shimura, T., Metabolic studies of the edematous cerebral cortex of the pyrithiamine-treated thiamine-deficient rat, 202
Tehovnik, E.J. and Yeomans, J.S., Contraversive circling elicited from the internal capsule and substantia nigra: evidence for a continuous axon bundle mediating circling, 269

Urbanski, H.F., see Ojeda, S.R., 339

Wada, H., see Sakata, T., 403
White, J.D., see La Gamma, E.F., 292
White, S.R., Penner, J.D., Speth, R.C. and Chan, J.Y.H., Angiotensin II receptors in the lumbar spinal cord of the rat, 195
Willis, W.D., see Harmann, P.A., 87
Wright, J.W., Quirk, W.S., Hanesworth, J.M. and Harding, J.W., Influence of aminopeptidase inhibitors on brain angiotensin metabolism and drinking in rats, 215

Yeomans, J.S., see Tehovnik, E.J., 269
Yoshimatsu, H., see Sakata, T., 403
Yoshino, Y., see Takahashi, H., 202